





Comparative Study of Juvenile American Shad Populations by Fin Ray and Scute Counts

by Paul R. Nichols



SPECIAL SCIENTIFIC REPORT-FISHERIES No. 525

UNITED STATES DEPARTMENT OF THE INTERIOR

FISH AND WILDLIFE SERVICE

BUREAU OF COMMERCIAL FISHERIES

UNITED STATES DEPARTMENT OF THE INTERIOR

Stewart L. Udall, *Secretary*

John A. Carver, Jr., *Under Secretary*

Stanley A. Cain, *Assistant Secretary for Fish and Wildlife and Parks*

FISH AND WILDLIFE SERVICE, Clarence F. Pautzke, *Commissioner*

BUREAU OF COMMERCIAL FISHERIES, Donald L. McKernan, *Director*

Comparative Study of Juvenile American Shad Populations by Fin Ray and Scute Counts

By

PAUL R. NICHOLS

United States Fish and Wildlife Service
Special Scientific Report--Fisheries No. 525

Washington, D.C.
February 1966

CONTENTS

Page

Introduction	1
Materials and methods	1
Analyses of meristic counts	2
North Atlantic area	2
Connecticut River	2
Hudson River	2
Comparison between Connecticut and Hudson Rivers	3
Chesapeake Bay area	4
Susquehanna River	4
Rappahannock River	4
York River	4
James River	5
Comparison between Chesapeake Bay tributaries	5
South Atlantic area	6
Neuse River	6
Edisto River	7
Ogeechee River	7
St. Johns River	7
Comparison between South Atlantic coast rivers	7
Relation between meristic counts and other factors	9
Summary	9
Acknowledgments	10
Literature cited	10

TABLES

1. Juvenile American shad collections, from 10 Atlantic coast rivers, examined in meristic studies	2
2. Frequencies of meristic counts from juvenile American shad in samples from the Connecticut River, Conn., and Mass	3
3. Frequencies of meristic counts from juvenile American shad in samples from the Hudson River, N.Y.	3
4. Analysis of variance on meristic counts for differences between locations and between years within rivers, and differences between rivers, for samples of juvenile American shad from the Connecticut and Hudson Rivers	4
5. Frequencies of meristic counts from juvenile American shad in samples from the Susquehanna River, Md.	4
6. Frequencies of meristic counts from juvenile American shad in samples from the Rappahannock River, Va.	4
7. Frequencies of meristic counts from juvenile American shad in samples from the York River tributaries, Va.	5
8. Frequencies of meristic counts from juvenile American shad in samples from the James River, Va.	5
9. Analysis of variance on meristic counts for differences between locations and between years within rivers, and differences between rivers, for samples of juvenile American shad from Chesapeake Bay tributaries	6
10. Frequencies of meristic counts from juvenile American shad in samples from the Neuse River, N.C.	6
11. Frequencies of meristic counts from juvenile American shad in samples from the Edisto River, S.C.	7
12. Frequencies of meristic counts from juvenile American shad in samples from the Ogeechee River, Ga.	7
13. Frequencies of meristic counts from juvenile American shad in samples from the St. Johns River, Fla.	8
14. Analysis of variance on meristic counts for differences between locations and between years within rivers, and differences between rivers, for samples of juvenile American shad from South Atlantic coast rivers.	8

Comparative Study of Juvenile American Shad Populations by Fin Ray and Scute Counts

By

PAUL R. NICHOLS, Fishery Biologist (Research)

Bureau of Commercial Fisheries Biological Laboratory
Beaufort, N.C.

ABSTRACT

Forty-five juvenile American shad, Alosa sapidissima (Wilson), collections, from 10 major shad producing rivers along the Atlantic coast of North America, were examined to see if differences in meristic counts suggested evidence of discrete river populations. Four meristic characters--pectoral, dorsal, and anal fin rays and scutes--were used. The difference in the counts between locations and between years within rivers was small compared to that between rivers. The differences in counts between rivers indicated that discrete populations of juvenile shad occurred in rivers.

INTRODUCTION

In studies to discover causes of the decline in yield of American shad, Alosa sapidissima (Wilson), and to determine factors favoring recovery of the fishery, it was essential that the number and distribution of populations be known.

The shad is widely distributed along the Atlantic coast from the St. Lawrence River, Canada, to the St. Johns River, Fla. This species is anadromous, spending most of its life in the ocean, but ascending coastal rivers to spawn. The spawning migrations into the rivers begin earliest in the southern part of the range (November in St. Johns River, Fla.) and are progressively later northward (June in St. John River, Canada). A female spawns about 250,000 eggs, and hatching occurs in 6 to 8 days at a water temperature of 17° C. The young shad stay in the rivers until autumn, attaining a length ranging from 75 to 145mm., and then migrate to sea. After spending from 2 to 6 years in the ocean, shad return to the rivers to spawn. Those spawning in rivers south of Cape Hatteras, N.C., normally die after spawning, while north of Cape Hatteras the proportion of fish spawning for the second time or more progressively increases from about 15 to 25 percent in Chesapeake Bay tributaries to about 45 to 55 percent in the Connecticut River.

Several workers have reported evidence of different shad populations along the Atlantic coast. Differences between shad from different areas based on meristic counts (Fischler, 1959; Hildebrand and Schroeder, 1928; Hill,

1959; Vladykov and Wallace, 1938), growth rates (Hammer, 1942;¹ Hildebrand and Schroeder, 1928), and fecundity (Davis, 1957; Lehman, 1953) indicated the occurrence of different populations. Recapture on the spawning ground of shad tagged in prior seasons indicated that they returned to their native streams to spawn (Hollis, 1948; Nichols, 1960). Also, the fact that the runs in the northern rivers were self-perpetuating and fluctuated independently (Talbot and Sykes, 1958) suggested different populations.

The purpose of this study was to determine if discrete populations of shad could be identified on the basis of consistent differences in counts of meristic characters in juveniles from 10 rivers. As used in this report, a "population" is a group of fish having similar meristic characteristics, of which the nature of origin, genotypic and/or phenotypic, of the characteristics has not been determined.

MATERIALS AND METHODS

For this study, the Atlantic coast was divided into three geographical areas: North Atlantic (Maine to Virginia); Chesapeake Bay (Maryland and Virginia); and South Atlantic (North Carolina to Florida). In 10 major shad producing rivers within these areas, 45 collections of juvenile shad, from 43 to 146 mm. fork length, were taken near spawning and

¹The homing instinct of the Chesapeake Bay shad, Alosa sapidissima (Wilson), as revealed by a study of their scales. Thesis (typewritten), 1942, University of Maryland, 45 p.

nursery areas (table 1). Fifty specimens were sampled at random from each collection, covering the size range in each, for the analysis.

Table 1.--Juvenile American shad collections, from 10 Atlantic coast rivers, examined in meristic studies

Area and river	Collecting gear	Locations	Collections	Specimens	Size range fork length
		Number	Number	Number	Mm.
North Atlantic:					
Connecticut...	Seine	2	6	300	55-146
Hudson.....	Seine	3	9	450	45- 82
Chesapeake Bay:					
Susquehanna...	Seine	1	1	50	48- 70
Rappahannock...	Trawl	2	2	100	64- 89
York.....	Trawl	2	6	300	54- 90
James.....	Trawl	2	3	150	55- 88
South Atlantic:					
Neuse.....	Seine	2	5	250	43-110
Edisto.....	Trawl	1	4	200	56- 83
Ogeechee.....	Trawl	2	4	200	54- 84
St. Johns....	Trawl	3	5	250	46- 74
		Total...	45	2,250	

In the North Atlantic area, collections of juvenile shad were taken from the Connecticut and Hudson Rivers. From the Connecticut, collections were taken above South Hadley Falls Dam at Holyoke, Mass., about 85 miles from the river mouth, and at Enfield, Conn., about 15 miles downstream of South Hadley Falls Dam, in the fall of 1954, 1957, and 1958. The collections taken above South Hadley Falls Dam were considered as an introduced population, because the dam blocked upstream migrating fish from this area for more than 100 years until a fish-passage facility was installed in 1952. In the Hudson, collections were taken at Piermont, N.Y., in the brackish water section about 30 miles from the river mouth; at Kingston Point, N.Y., in the freshwater section about 50 miles upstream from Piermont; and at Catskill, N.Y., about 25 miles upstream from Kingston Point, in the autumn of 1950 and 1951. Additional collections were taken at Kingston Point in 1954, 1957, and 1958.

In the Chesapeake Bay area, collections of juvenile shad were taken from the Susquehanna, Rappahannock, York, and James Rivers. From the Susquehanna River, collections were taken below the Conowingo Dam in 1958; from the Rappahannock River, at Long Point in 1954 and in Batchelors Bay in 1958; from the York River, at the Pamunkey Indian Reservation in 1953, 1954, 1956, and 1958 and at the Mattaponi Indian Reservation in 1954 and 1958; and from the James River, at Walcot Wharf, Va., in 1954 and at Claremont Beach, Va., in 1954 and 1958.

In the South Atlantic area, collections of juvenile shad were taken from the Neuse, Edisto, Ogeechee, and St. Johns Rivers. From the Neuse River, collections were taken at Bridgeton, N.C., in 1950, 1954, 1957, and 1958

and at Streets Ferry, N.C., in 1954; collections from the Edisto River at Crosby Landing, S.C., were available for 1938 and 1939 and were taken in 1957 and 1958; collections from the Ogeechee River were available for 1938 and 1939 from Kings Ferry, Ga., and were taken in 1957 and 1958 at the State Park, near Richmond Hill, Ga.; and collections from the St. Johns River were taken at Mandarin, Fla., in 1954, in Lake Harney in 1954, and at Palatka, Fla., in 1954, 1957, and 1958.

Using a binocular microscope, counts were made of left pectoral, dorsal, and anal fin rays and total scutes. Fin ray counts included all rudiments, and the last elements in anal and dorsal fins, originating from the same base, were counted as one ray. The dorsal fin origin often required dissection to expose embedded rays. Scales occasionally had to be removed to expose enveloped scutes and anal fin rays. No attempt was made to separate scutes into anterior and posterior counts. Not a single abnormal fin or scute was encountered out of the 2,250 specimens examined.

Analysis of variance (Snedecor, 1956; Steel and Torrie, 1960) was used to test if meristic count means of specimens were statistically different at the 1 percent level (indicated by two asterisks in the tables) between rivers, locations within rivers, and years within rivers. Before comparing the means, group variances were tested for homogeneity.

ANALYSES OF MERISTIC COUNTS

Differences and similarities in meristic counts for samples of shad from within individual rivers and between rivers are discussed in the following sections by geographical area.

North Atlantic Area

Meristic counts were made of juvenile shad taken at each location in the Connecticut and Hudson Rivers (tables 2 and 3).

Connecticut River.--Mean meristic counts for the Holyoke samples (above Hadley Falls Dam) in most instances were slightly higher than those for Enfield samples (below Hadley Falls Dam). The difference in pectoral fin ray counts was significant between locations (table 4). The difference in each mean meristic count was nonsignificant between years for the Enfield samples. No analysis was made for differences between years in the meristic counts from above Hadley Falls Dam.

Hudson River.--Differences in meristic counts were not significant between locations (Piermont-Kingston-Catskill) for the years 1950 and 1951. Since Kingston Point was the

Table 2.--Frequencies of meristic counts from juvenile American shad in samples from the Connecticut River, Conn., and Mass.

Location	Year	Number of pectoral fin rays										Mean	Standard deviation
		13	14	15	16	17	18	19	20	21	22		
<u>Frequency</u>													
Enfield, Conn.	1954	3	30	17	-	-	-	-	-	-	-	15.28	0.5729
Do.	1957	7	29	14	-	-	-	-	-	-	-	15.14	0.6392
Do.	1958	8	23	19	-	-	-	-	-	-	-	15.22	0.7083
Holyoke, Mass.	1954	-	12	33	4	1	-	-	-	-	-	15.88	0.6273
Do.	1957	-	16	28	6	-	-	-	-	-	-	15.80	0.6389
Do.	1958	-	13	33	4	-	-	-	-	-	-	15.82	0.5602
<u>Number of dorsal fin rays</u>													
<u>15 16 17 18 19 20 21 22</u>													
<u>Frequency</u>													
Enfield, Conn.	1954	-	6	30	14	-	-	-	-	-	-	18.16	0.6181
Do.	1957	-	6	33	11	-	-	-	-	-	-	18.10	0.5803
Do.	1958	-	4	33	13	-	-	-	-	-	-	18.18	0.5602
Holyoke, Mass.	1954	-	4	34	11	-	-	-	-	-	-	18.18	0.5956
Do.	1957	-	1	31	18	-	-	-	-	-	-	18.34	0.5794
Do.	1958	2	10	20	17	-	-	-	-	-	-	18.10	0.8864
<u>Number of anal fin rays</u>													
<u>18 19 20 21 22 23 24 25</u>													
<u>Frequency</u>													
Enfield, Conn.	1954	-	8	29	12	1	-	-	-	-	-	21.12	0.6893
Do.	1957	1	8	23	9	9	-	-	-	-	-	21.34	1.0224
Do.	1958	4	11	16	13	6	-	-	-	-	-	21.12	1.1364
Holyoke, Mass.	1954	1	6	25	18	-	-	-	-	-	-	21.20	0.7284
Do.	1957	1	5	27	16	1	-	-	-	-	-	21.22	0.7365
Do.	1958	2	10	19	16	2	1	-	-	-	-	21.18	1.0039
<u>Number of scutes</u>													
<u>33 34 35 36 37 38 39 40</u>													
<u>Frequency</u>													
Enfield, Conn.	1954	-	5	21	20	3	1	-	-	-	-	36.48	0.8389
Do.	1957	1	4	15	19	11	-	-	-	-	-	36.70	0.9742
Do.	1958	-	4	12	22	12	-	-	-	-	-	36.84	0.8889
Holyoke, Mass.	1954	1	2	15	23	8	1	-	-	-	-	36.76	0.9161
Do.	1957	-	3	18	21	8	-	-	-	-	-	36.68	0.8192
Do.	1958	-	6	10	16	16	2	-	-	-	-	36.96	1.0872

only location that had a sufficient number of data for comparing different young, samples taken at Kingston Point only were used to test for differences between years. There were no significant differences between years (1950, 1951, 1954, 1957, 1958) for these samples (table 4).

Comparison between Connecticut and Hudson Rivers.--Differences in the meristic counts for the Hudson and Connecticut Rivers samples were tested for significance only for the years for which collections were available from both rivers (1954, 1957, 1958). Therefore, only the collections taken at Kingston Point on the Hudson and at Enfield on the Connecticut were used in the comparisons. The Enfield samples were considered representative of the Connecticut River popula-

Table 3.--Frequencies of meristic counts from juvenile American shad in samples from the Hudson River, N. Y.

Location	Year	Number of pectoral fin rays										Mean	Standard deviation
		13	14	15	16	17	18	19	20	21			
		<u>Frequency</u>											
Piermont	1950	4	20	23	3						15.50	0.7354	
Do.	1951	-	23	24	3						15.60	0.6061	
Kingston Point	1950	2	15	28	5						15.72	0.7010	
Do.	1951	-	19	29	2						15.66	0.5573	
Do.	1954	1	19	28	2						15.62	0.6024	
Do.	1957	-	10	34	6						15.92	0.5657	
Do.	1958	1	6	36	7						15.98	0.5887	
Catskill	1950	3	17	27	3						15.58	0.6999	
Do.	1951	1	27	21	1						15.28	0.5771	
		<u>Number of dorsal fin rays</u>											
		16	17	18	19	20	21	22	23	24			
		<u>Frequency</u>											
Piermont	1950	3	28	13	6						18.44	0.7866	
Do.	1951	1	25	20	4						18.54	0.6764	
Kingston Point	1950	2	24	20	4						18.52	0.7068	
Do.	1951	4	18	27	1						18.50	0.6776	
Do.	1954	3	27	20	-						18.34	0.5928	
Do.	1957	3	24	22	1						18.42	0.6417	
Do.	1958	3	24	21	2						18.44	0.6749	
Catskill	1950	6	20	21	3						18.24	0.7440	
Do.	1951	4	19	26	1						18.48	0.6773	
		<u>Number of anal fin rays</u>											
		18	19	20	21	22	23	24	25	26			
		<u>Frequency</u>											
Piermont	1950	1	5	20	21	3	-				21.40	0.8330	
Do.	1951	-	4	12	27	4	3				21.80	0.9258	
Kingston Point	1950	-	7	21	16	4	2				21.46	0.9733	
Do.	1951	-	6	19	17	7	1				21.56	0.9510	
Do.	1954	-	10	21	13	6	-				21.30	0.9313	
Do.	1957	2	12	26	8	2	-				20.92	0.8533	
Do.	1958	-	10	29	9	2	-				21.06	0.7398	
Catskill	1950	2	7	17	18	6	-				21.36	1.0079	
Do.	1951	1	12	20	13	4	-				21.10	0.9478	
		<u>Number of scutes</u>											
		33	34	35	36	37	38	39	40	41			
		<u>Frequency</u>											
Piermont	1950	-	3	4	17	19	6	1			37.50	1.0736	
Do.	1951	1	1	7	21	11	9	-			37.34	1.1178	
Kingston Point	1950	-	11	15	20	3	1	-			37.36	0.9638	
Do.	1951	-	1	10	19	14	6	-			37.28	0.9906	
Do.	1954	1	2	4	13	20	8	2			37.62	1.1952	
Do.	1957	-	1	7	20	15	5	2			37.44	1.0529	
Do.	1958	-	4	3	15	26	2	-			37.38	0.9666	
Catskill	1950	-	1	5	16	20	7	1			37.58	0.9897	
Do.	1951	-	3	7	25	10	5	-			37.00	0.8562	

tion, while the Kingston Point sample was considered representative of the Hudson River population.

The mean meristic counts for Hudson River shad generally were higher than those of Connecticut River fish. Significant differences were found between rivers for all meristic counts, except anal fin rays (table 4). The interaction between years and rivers was not significant. The significant difference in three of the four counts indicated that discrete populations occurred in each river.

Table 4.--Analysis of variance on meristic counts for differences between locations and between years within rivers, and differences between rivers, for samples of juvenile American shad from the Connecticut and Hudson Rivers

River	Component	Degrees of freedom (n ₁ , n ₂)	F-value
Connecticut	Between locations:		
	Pectoral fin rays.....	1,296	73.923**
	Dorsal fin rays.....	1,296	0.680
	Anal fin rays.....	1,296	0.060
	Scutes.....	1,296	1.403
	Between years:		
	(Enfield Dam only)		
	Pectoral fin rays.....	2,147	0.605
	Dorsal fin rays.....	2,147	0.291
	Anal fin rays.....	2,147	0.854
	Scutes.....	2,147	2.027
Hudson	Between locations:		
	Pectoral fin rays.....	2,296	1.955
	Dorsal fin rays.....	2,296	1.317
	Anal fin rays.....	2,296	3.455
	Scutes.....	2,296	0.262
	Between years:		
	(Kingston Point only)		
	Pectoral fin rays.....	4,245	3.395
	Dorsal fin rays.....	4,245	0.496
	Anal fin rays.....	4,245	4.494
	Scutes.....	4,245	0.665
Connecticut and Hudson	Between rivers:		
	Pectoral fin rays.....	1,296	76.506**
	Dorsal fin rays.....	1,296	12.853**
	Anal fin rays.....	1,296	0.915
	Scutes.....	1,296	48.726**

** Statistically different at 1 percent level.

Chesapeake Bay Area

Frequencies of the meristic counts for the samples of juvenile shad from the Chesapeake Bay tributaries are given in tables 5, 6, 7, and 8.

Susquehanna River.--Only one sample of juvenile shad was available from the Susquehanna River, collected in 1958 below Conowingo Dam (table 5), so no comparisons could be made.

Rappahannock River.--Mean meristic counts for the Rappahannock River samples were similar, and tests for differences in the counts between years were not significant (table 6). No comparisons were made, because the only sample taken in each of the two years was taken at different locations.

York River.--Mean meristic counts for the York River samples were similar between years and locations (table 7). Based on the

Table 5.--Frequencies of meristic counts from juvenile American shad in samples from the Susquehanna River, Md.

Location	Year	Number of pectoral fin rays								Mean	Standard deviation
		12	13	14	15	16	17	18	19		
Conowingo Dam	1958	Frequency								16.12	0.5206
		- 4 36 10									
		Number of dorsal fin rays									
		15 16 17 18 19 20 21 22									
Conowingo Dam	1958	Frequency								18.34	0.4785
		- 33 17 -									
		Number of anal fin rays									
		18 19 20 21 22 23 24 25									
Conowingo Dam	1958	Frequency								21.28	0.7570
		- 8 21 20 1 -									
		Number of scutes									
		33 34 35 36 37 38 39 40									
Conowingo Dam	1958	Frequency								37.22	0.7083
		- - 7 26 16 1									

Table 6.--Frequencies of meristic counts from juvenile American shad in samples from the Rappahannock River, Va.

Location	Year	Number of pectoral fin rays										Mean	Standard deviation
		11	12	13	14	15	16	17	18	19	20		
Long Point Batchelors Bay	1954 1958	Frequency										15.25 15.54	0.6713 0.5789
		Number of dorsal fin rays											
		14	15	16	17	18	19	20	21	22	23		
Long Point Batchelors Bay	1954 1958	Frequency										18.54 18.54	0.7060 0.7060
		Number of anal fin rays											
		17	18	19	20	21	22	23	24	25	26		
Long Point Batchelors Bay	1954 1958	Frequency										21.64 21.38	0.7559 0.9666
		Number of scutes											
		32	33	34	35	36	37	38	39	40	41		
Long Point Batchelors Bay	1954 1958	Frequency										36.42 36.12	0.7848 1.0812

Table 7.--Frequencies of meristic counts from juvenile American shad in samples from the York River tributaries, Va.

Location	Year	Number of pectoral fin rays								Mean	Standard deviation
		12	13	14	15	16	17	18	19		
		<u>Frequency</u>									
Pamunkey River	1953	2	31	17	-					16.30	0.5440
Do.	1954	2	24	23	1					16.46	0.6131
Do.	1956	4	33	12	1					16.20	0.6061
Do.	1958	-	28	22	-					16.44	0.5014
Mattaponi River	1954	2	26	21	1					16.42	0.6091
Do.	1958	4	24	22	-					16.36	0.6312
		<u>Number of dorsal fin rays</u>									
		15	16	17	18	19	20	21	22		
		<u>Frequency</u>									
Pamunkey River	1953	-	24	25	1					18.54	0.5425
Do.	1954	-	26	22	2					18.52	0.5799
Do.	1956	1	21	25	3					18.60	0.6389
Do.	1958	3	15	27	5					18.68	0.7407
Mattaponi River	1954	1	18	21	10					18.52	0.5799
Do.	1958	-	26	22	2					18.80	0.7825
		<u>Number of anal fin rays</u>									
		18	19	20	21	22	23	24	25		
		<u>Frequency</u>									
Pamunkey River	1953	1	7	27	12	3				21.18	0.8254
Do.	1954	-	10	18	20	2				21.28	0.8340
Do.	1956	-	7	22	16	5				21.38	0.8545
Do.	1958	-	12	16	20	2				21.24	0.8404
Mattaponi River	1954	-	11	28	10	1				21.02	0.7140
Do.	1958	2	14	23	9	2				20.90	0.8864
		<u>Number of scutes</u>									
		33	34	35	36	37	38	39	40		
		<u>Frequency</u>									
Pamunkey River	1953	1	3	14	21	10	1			36.78	0.9750
Do.	1954	-	3	14	21	9	3			36.94	0.9742
Do.	1956	2	3	8	18	16	3			37.04	1.1599
Do.	1958	-	1	9	28	11	1			37.04	0.7548
Mattaponi River	1954	2	1	12	22	11	2			36.90	1.0351
Do.	1958	-	2	11	22	13	2			37.04	0.9026

samples from 1954 and 1958 only, differences in the meristic counts were not significant between locations. Based on the Pamunkey River samples only, differences between years also were not significant (table 9).

James River.--Mean meristic counts for the James River samples were similar between years and between locations (table 8). Based on the 1954 samples, counts were not significant between the two locations, and for the

Table 8.--Frequencies of meristic counts from juvenile American shad in samples from the James River, Va.

Location	Year	Number of pectoral fin rays								Mean	Standard deviation
		12	13	14	15	16	17	18	19		
		<u>Frequency</u>									
Walcot Wharf	1954	-	21	28	1					15.60	0.5345
Claremont Beach	1954	1	19	27	3					15.64	0.6312
Do.	1958	-	12	34	4					15.84	0.5481
		<u>Number of dorsal fin rays</u>									
		15	16	17	18	19	20	21	22		
		<u>Frequency</u>									
Walcot Wharf	1954	1	29	20	-					18.38	0.5303
Claremont Beach	1954	5	29	16	-					18.22	0.6158
Do.	1958	5	32	11	2					18.22	0.7083
		<u>Number of anal fin rays</u>									
		18	19	20	21	22	23	24	25		
		<u>Frequency</u>									
Walcot Wharf	1954	17	21	9	3					20.70	0.8631
Claremont Beach	1954	24	21	5	-					20.62	0.6667
Do.	1958	13	25	11	1					21.00	0.7559
		<u>Number of scutes</u>									
		33	34	35	36	37	38	39	40		
		<u>Frequency</u>									
Walcot Wharf	1954	2	6	27	12	2	1			36.18	0.9190
Claremont Beach	1954	1	8	20	14	7	-			36.36	0.9848
Do.	1958	2	6	20	18	4	-			36.32	0.9355

Claremont Beach samples the differences between years were not significant.

Comparison between Chesapeake Bay tributaries.--Data for all locations and all years for which collections were available were combined in testing for significant differences in meristic counts between two rivers. There was a significant difference in the meristic counts between the York and James fish in all instances; between the Rappahannock and James, the Rappahannock and York, and the James and Susquehanna in three instances; and between the Rappahannock and Susquehanna, and the York and Susquehanna in two instances (table 9). Where applicable, the interaction between years and rivers was not significant. The differences in meristic counts between rivers indicated that discrete populations of shad occurred in the Chesapeake Bay tributaries.

Table 9.--Analysis of variance on meristic counts for differences between locations and between years within rivers, and differences between rivers, for samples of juvenile American shad from Chesapeake Bay tributaries

River	Component	Degrees of freedom (n ₁ , n ₂)	F-value
York	Between locations:		
	Pectoral fin rays.....	1,197	0.514
	Dorsal fin rays.....	1,197	0.391
	Anal fin rays.....	1,197	6.569
	Scutes.....	1,197	0.000
	Between years:		
	(Pamunkey River only)		
	Pectoral fin rays.....	3,196	2.335
	Dorsal fin rays.....	3,196	0.651
	Anal fin rays.....	3,196	0.493
	Scutes.....	3,196	0.825
James	Between locations:		
	Pectoral fin rays.....	1, 98	0.117
	Dorsal fin rays.....	1, 98	1.938
	Anal fin rays.....	1, 98	0.269
	Scutes.....	1, 98	0.893
	Between years:		
	(Claremont Beach only)		
	Pectoral fin rays.....	1, 98	4.914
	Dorsal fin rays.....	1, 98	1.635
	Anal fin rays.....	1, 98	3.419
	Scutes.....	1, 98	0.570
Rappahannock-James	Between rivers:		
	Pectoral fin rays.....	1,248	13.258**
	Dorsal fin rays.....	1,248	9.922**
	Anal fin rays.....	1,248	46.227**
	Scutes.....	1,248	0.011
Rappahannock-York	Between rivers:		
	Pectoral fin rays.....	1,398	188.925**
	Dorsal fin rays.....	1,398	0.835
	Anal fin rays.....	1,398	10.885**
	Scutes.....	1,398	37.095**
Rappahannock-Susquehanna	Between rivers:		
	Pectoral fin rays.....	1,148	46.512**
	Dorsal fin rays.....	1,148	3.287
	Anal fin rays.....	1,148	2.111
	Scutes.....	1,148	38.965**
James-York	Between rivers:		
	Pectoral fin rays.....	1,448	131.156**
	Dorsal fin rays.....	1,448	27.412**
	Anal fin rays.....	1,448	22.951**
	Scutes.....	1,448	47.494**
James-Susquehanna	Between rivers:		
	Pectoral fin rays.....	1,198	21.394**
	Dorsal fin rays.....	1,198	0.476
	Anal fin rays.....	1,198	16.101**
	Scutes.....	1,198	41.121**
York-Susquehanna	Between rivers:		
	Pectoral fin rays.....	1,348	7.570**
	Dorsal fin rays.....	1,348	7.859**
	Anal fin rays.....	1,348	0.798
	Scutes.....	1,348	3.550

** Statistically different at 1 percent level.

South Atlantic Area

Meristic counts were made of juvenile shad taken for certain years at locations in the Neuse, Edisto, Ogeechee, and St. Johns Rivers of the South Atlantic area (tables 10, 11, 12, and 13).

Neuse River.--Based on the 1954 samples (table 10), differences in the meristic counts were not significant between the Bridgeton and Streets Ferry samples. Differences in the meristic counts were not significant between years for the Bridgeton samples (table 14).

Table 10.--Frequencies of meristic counts from juvenile American shad in samples from the Neuse River, N. C.

Location	Year	Number of pectoral fin rays								Mean	Standard deviation
		12	13	14	15	16	17	18	19		
<u>Frequency</u>											
Bridgeton	1950	-	9	36	5	-				15.92	0.5284
Do.	1954	-	12	31	6	1				15.92	0.6652
Do.	1957	1	7	28	14	-				16.14	0.7001
Do.	1958	-	7	25	14	-				16.22	0.6788
Streets Ferry	1954	1	14	29	6	-				15.80	0.6701
<u>Number of dorsal fin rays</u>											
<u>15 16 17 18 19 20 21 22</u>											
<u>Frequency</u>											
Bridgeton	1950	3	17	26	4	-				18.56	0.7329
Do.	1954	-	10	33	7	-				18.92	0.5859
Do.	1957	-	16	28	6	-				18.80	0.6389
Do.	1958	-	21	25	3	1				18.68	0.6833
Streets Ferry	1954	-	10	33	7	-				18.74	0.8033
<u>Number of anal fin rays</u>											
<u>18 19 20 21 22 23 24 25</u>											
<u>Frequency</u>											
Bridgeton	1950	2	8	22	16	2	-			21.56	0.8889
Do.	1954	-	4	15	20	10	1			21.58	0.9708
Do.	1957	2	6	15	21	6	-			21.46	0.9941
Do.	1958	2	5	20	18	5	-			21.38	0.9452
Streets Ferry	1954	-	6	14	21	9	-			21.78	0.9322
<u>Number of scutes</u>											
<u>33 34 35 36 37 38 39 40</u>											
<u>Frequency</u>											
Bridgeton	1950	1	7	17	15	7	3			36.58	1.1445
Do.	1954	2	8	20	16	4	-			35.24	0.9596
Do.	1957	-	17	18	8	7	-			36.10	1.0351
Do.	1958	-	3	27	14	5	1			36.48	0.8389
Streets Ferry	1954	4	5	17	18	6	-			36.04	1.0806

Edisto River.--Samples were collected from only one location in the Edisto River (table 11). The differences in the counts between years were not significant (table 14).

Ogeechee River.--Since samples were obtained in different years from the two locations in the Ogeechee River, each location was analyzed separately in testing for differences in meristic counts between years, and no comparisons were made between locations (table 12). The differences in the counts between years were not significant (table 14).

Table 11.--Frequencies of meristic counts from juvenile American shad in samples from the Edisto River, S. C.

Location	Year	Number of										Mean	Standard deviation
		pectoral fin rays											
		12	13	14	15	16	17	18	19				
		<u>Frequency</u>											
Crosby Landing	1938				6	28	16	-			16.20	0.6389	
Do.	1939				4	27	18	1			16.32	0.6528	
Do.	1957				1	31	17	1			16.36	0.5628	
Do.	1958				1	35	13	1			16.28	0.5360	
		<u>Number of</u>											
		<u>dorsal fin rays</u>											
		15	16	17	18	19	20	21	22				
		<u>Frequency</u>											
Crosby Landing	1938				2	20	25	3			18.58	0.6728	
Do.	1939				2	24	22	2			18.48	0.6465	
Do.	1957				1	23	24	2			18.54	0.6131	
Do.	1958				-	20	28	2			18.64	0.5628	
		<u>Number of</u>											
		<u>anal fin rays</u>											
		18	19	20	21	22	23	24	25				
		<u>Frequency</u>											
Crosby Landing	1938				2	15	22	10	1		20.86	0.8574	
Do.	1939				2	20	17	11	-		20.74	0.8526	
Do.	1957				2	23	18	7	-		20.60	0.7825	
Do.	1958				3	15	24	8	-		20.74	0.8033	
		<u>Number of scutes</u>											
		33	34	35	36	37	38	39	40				
		<u>Frequency</u>											
Crosby Landing	1938				1	3	22	17	6	1	36.54	0.9304	
Do.	1939				-	4	24	17	3	2	36.50	0.8864	
Do.	1957				-	7	24	15	2	2	36.36	0.9205	
Do.	1958				1	7	20	18	4	-	36.34	0.8947	

Table 12.--Frequencies of meristic counts from juvenile American shad in samples from the Ogeechee River, Ga.

Location	Year	Number of pectoral fin rays										Mean	Standard deviation
		12	13	14	15	16	17	18	19				
		<u>Frequency</u>											
Kings Ferry	1938				5	36	9	-				16.30	0.6145
Do.	1939				3	30	16	1				16.07	0.5284
State Park	1957				1	39	10	1				16.18	0.4375
Do.	1958				-	34	15	1				16.34	0.5194
		<u>Number of dorsal fin rays</u>											
		15	16	17	18	19	20	21	22				
		<u>Frequency</u>											
Kings Ferry	1938				13	34	3	-				18.86	0.6704
Do.	1939				15	27	8	-				19.00	0.5345
State Park	1957				13	32	4	1				18.86	0.6392
Do.	1958				13	30	6	1				18.90	0.6776
		<u>Number of anal fin rays</u>											
		18	19	20	21	22	23	24	25				
		<u>Frequency</u>											
Kings Ferry	1938		2	13	21	11	3					21.00	0.9476
Do.	1939		2	8	26	11	3					21.10	0.8864
State Park	1957		3	8	21	16	2					21.12	0.9398
Do.	1958		1	11	23	7	8					21.20	1.0302
		<u>Number of scutes</u>											
		33	34	35	36	37	38	39	40				
		<u>Frequency</u>											
Kings Ferry	1938		2	2	24	17	5	-				36.64	1.1205
Do.	1939		-	9	13	18	7	3				36.43	1.0529
State Park	1957		-	9	17	18	6	-				36.42	0.9278
Do.	1958		-	6	19	17	8	-				36.54	0.9082

St. Johns River.--Based on the 1954 samples from the St. Johns River (table 13), differences in the meristic counts were not significant between the three locations. Differences between years were tested for the Palatka samples only and were not significant (table 14).

Comparison between South Atlantic coast rivers.--Data for all locations and all years for which collections were available were combined in testing for significant differences in meristic counts between two rivers. There was

Table 13.--Frequencies of meristic counts from juvenile American shad in samples from the St. Johns River, Fla.

Location	Year	Number of pectoral fin rays								Mean	Standard deviation		
		12	13	14	15	16	17	18	19				
		<u>Frequency</u>											
Mandarin	1954					1	32	17		16.32	0.4518		
Lake Harney	1954					7	28	15		16.16	0.6503		
Palatka	1954					1	38	11		16.20	0.5127		
Do.	1957					6	27	17		16.22	0.6481		
Do.	1958					5	37	8		16.06	0.5115		
		<u>Number of dorsal fin rays</u>											
		<u>15 16 17 18 19 20 21 22</u>											
		<u>Frequency</u>											
Mandarin	1954					12	29	9	-	18.94	0.6389		
Lake Harney	1954					11	24	14	1	19.10	0.7626		
Palatka	1954					10	30	10	-	19.00	0.6518		
Do.	1957					10	33	7	-	18.94	0.5859		
Do.	1958					11	29	10	-	18.94	0.6543		
		<u>Number of anal fin rays</u>											
		<u>18 19 20 21 22 23 24 25</u>											
		<u>Frequency</u>											
Mandarin	1954					5	19	18	8	21.58	0.8053		
Lake Harney	1954					2	21	21	5	21.64	0.8020		
Palatka	1954					3	19	23	4	21.62	0.8827		
Do.	1957					5	21	19	4	21.50	0.8631		
Do.	1958					2	14	28	6	21.76	0.7160		
		<u>Number of scutes</u>											
		<u>33 34 35 36 37 38 39 40</u>											
		<u>Frequency</u>											
Mandarin	1954					-	3	11	26	10	36.86	0.9313	
Lake Harney	1954					-	6	11	15	14	4	36.98	1.1516
Palatka	1954					-	4	17	21	6	2	36.70	0.8084
Do.	1957					1	6	20	19	4	-	36.38	0.8781
Do.	1958					-	4	20	23	3	-	36.50	0.7354

a significant difference in the meristic counts between the Edisto and Neuse, the St. Johns and Neuse, and the St. Johns and Edisto in three

Table 14.--Analysis of variance on meristic counts for differences between locations and between years within rivers, and differences between rivers, for samples of juvenile American shad from South Atlantic coast rivers

River	Component	Degrees of freedom (n ₁ , n ₂)	F-value
Neuse	Between locations:		
	Pectoral fin rays.....	1, 98	0.808
	Dorsal fin rays.....	1, 98	2.023
	Anal fin rays.....	1, 98	1.104
	Scutes.....	1, 98	0.239
	Between years:		
	(Bridgeton only)		
	Pectoral fin rays.....	3, 196	2.822
	Dorsal fin rays.....	3, 196	3.019
	Anal fin rays.....	3, 196	1.732
Edisto	Between years:		
	(State Park only)		
	Pectoral fin rays.....	3, 196	0.649
	Dorsal fin rays.....	3, 196	0.580
	Anal fin rays.....	3, 196	0.831
	Scutes.....	3, 196	0.606
Ogeechee	Between years:		
	(State Park only)		
	Pectoral fin rays.....	1, 98	2.775
	Dorsal fin rays.....	1, 98	0.092
	Anal fin rays.....	1, 98	0.165
	Scutes.....	1, 98	0.427
	Between years:		
	(Kings Ferry only)		
	Pectoral fin rays.....	1, 98	3.685
St. Johns	Between years:		
	(Kings Ferry only)		
	Pectoral fin rays.....	1, 98	0.245
	Dorsal fin rays.....	1, 98	0.297
	Anal fin rays.....	1, 98	0.846
	Scutes.....	1, 98	0.846
	Between locations:		
	Pectoral fin rays.....	2, 147	1.170
	Dorsal fin rays.....	2, 147	0.693
	Anal fin rays.....	2, 147	0.068
	Scutes.....	2, 147	1.043

instances; between the Ogeechee and Neuse, and the Ogeechee and Edisto in two instances; and between the St. Johns and Ogeechee in one instance (table 14). These differences indicated that discrete populations of shad occurred in the South Atlantic coast rivers.

Table 14. --Analysis of variance on meristic counts for differences between locations and between years within rivers, and differences between rivers, for samples of juvenile American shad from South Atlantic coast rivers--Continued

River	Component	Degrees of freedom (n_1, n_2)	F-value
	Between years: (Palatka only)		
	Pectoral fin rays.....	2,147	2.732
	Dorsal fin rays.....	2,147	0.068
	Anal fin rays.....	2,147	1.306
	Scutes.....	2,147	4.763
Edisto-Neuse	Between rivers:		
	Pectoral fin rays.....	1,448	23.119**
	Dorsal fin rays.....	1,448	8.469**
	Anal fin rays.....	1,448	74.019**
	Scutes.....	1,448	0.898
Ogeechee-Neuse	Between rivers:		
	Pectoral fin rays.....	1,448	15.087**
	Dorsal fin rays.....	1,448	3.040
	Anal fin rays.....	1,448	16.382**
	Scutes.....	1,448	2.839
Ogeechee-Edisto	Between rivers:		
	Pectoral fin rays.....	1,398	1.305
	Dorsal fin rays.....	1,398	22.185**
	Anal fin rays.....	1,398	17.366**
	Scutes.....	1,398	0.614
St. Johns-Neuse	Between rivers:		
	Pectoral fin rays.....	1,498	12.160**
	Dorsal fin rays.....	1,498	16.679**
	Anal fin rays.....	1,498	3.465
	Scutes.....	1,498	14.738**
St. Johns-Edisto	Between rivers:		
	Pectoral fin rays.....	1,448	3.196
	Dorsal fin rays.....	1,448	50.127**
	Anal fin rays.....	1,448	130.011**
	Scutes.....	1,448	8.140**
St. Johns-Ogeechee	Between rivers:		
	Pectoral fin rays.....	1,448	0.396
	Dorsal fin rays.....	1,448	5.013
	Anal fin rays.....	1,448	38.413**
	Scutes.....	1,448	3.640

** Statistically different at 1 percent level.

RELATION BETWEEN MERISTIC COUNTS AND OTHER FACTORS

In certain instances significant differences were found in meristic counts between young shad from neighboring streams and among streams within large geographical areas. Whether the cause of these differences was primarily genetic or was environmental variation under which the fish developed, or a combination of both, would not affect the findings. Physical and chemical data on the environment at spawning time were not available, so the relation between differences or shifts in meristic counts between two separate rivers and environmental factors was not known. Lindsey (1957), Raney and Woolcott (1955), and Tåning (1952), and others, although working on species other than shad, listed temperature as the obvious environmental factor that may produce differences at the time these characters are formed in the embryo. Although shad spawn earlier in southern rivers and progressively later in northern rivers, widely separated populations spawn and eggs and larvae develop under about the same water temperature range. Laboratory rearing of the

species under controlled conditions would be needed to demonstrate clearly the relation between meristic counts and temperature.

There was no consistent latitudinal cline in the meristic counts examined over the entire geographic range sampled. Fin ray counts were higher in southern rivers and lower in northern rivers, with intermediate counts in between. Scute counts were higher in northern rivers and lower in southern rivers. The Connecticut River samples had the lowest mean count in pectoral and dorsal fin rays, and the St. Johns River samples had the highest mean counts in dorsal and anal fin rays. However, the mean scute counts, which showed the greatest difference, reversed this pattern with the high in the Hudson River being more than one unit larger than the low counts in the Neuse River.

There were slight variations in meristic counts on juvenile shad between years and locations within a river, but these were not significant and were small compared to the differences in the counts between rivers. The differences in meristic counts between rivers indicated that discrete juvenile populations exist. For a better understanding of shad populations, future work should include studies of the relation between the meristic counts of juvenile shad and environmental variations under which the fish developed.

SUMMARY

To test if shad populations in various rivers could be separated by differences in meristic count, collections of juvenile shad from 10 rivers along the Atlantic coast were compared by analysis of variance. Counts of pectoral, dorsal, and anal fin rays and scutes were used.

Differences in counts of meristic characteristics within individual rivers and between rivers within geographical areas were as follows:

1. Within individual rivers, no significant differences were found between locations and between years except between locations for pectoral fin rays in the Connecticut River.

2. In the North Atlantic area, significant differences were found between the Hudson and Connecticut Rivers for all counts except anal fin rays.

3. In Chesapeake Bay tributaries, significant differences were found between the York and James Rivers for all counts; between the Rappahannock and James Rivers, the Rappahannock and York Rivers, and the James and Susquehanna Rivers for three of the counts; and between the Rappahannock and Susquehanna, and the York and Susquehanna Rivers for two of the counts.

4. In the South Atlantic area, significant differences were found between the Neuse and Edisto Rivers, and the Neuse and St. Johns

Rivers for three of the counts; between the Neuse and Ogeechee Rivers, and the Ogeechee and Edisto Rivers for two of the counts; and between the Ogeechee and St. Johns Rivers for one of the counts.

These findings indicated that discrete populations of shad occurred in Atlantic coast rivers.

ACKNOWLEDGMENTS

William H. Massmann² provided collections of juvenile shad from the York River taken in 1954 and 1956, and Richard R. Whitney³ gave collections from the Susquehanna River taken in 1958. Edward C. Raney of Cornell University, Ithaca, N.Y., loaned collections from the Rappahannock and James Rivers taken in 1954.

LITERATURE CITED

DAVIS, WILLIAM S.

1957. Ova production of American shad in Atlantic coast rivers. U.S. Fish Wildl. Serv., Res. Rep. 49, 5 p.

FISCHLER, KENNETH J.

1959. Contributions of Hudson and Connecticut Rivers to New York-New Jersey shad catches of 1956. U.S. Fish Wildl. Serv., Fish. Bull. 60:161-174.

HILDEBRAND, SAMUEL F., and WILLIAM C. SCHROEDER.

1928. Fishes of the Chesapeake Bay. Bull. U.S. Bur. Fish. 43(pt. 1):1-366.

HILL, DONALD R.

1959. Some uses of statistical analysis in classifying races of American shad (Alosa sapidissima). U.S. Fish Wildl. Serv., Fish. Bull. 59:269-286.

HOLLIS, EDGAR H.

1948. The homing tendency of shad. Science 108(2804):332-333.

²Now with Sport Fishing Institute, Bond Bldg., Washington, D.C.

³Now with Bureau of Commercial Fisheries Biological Laboratory, P.O. Box 6317, Point Loma Station, San Diego, Calif.

LEACH, GLEN C.

1925. Artificial propagation of shad. [U.S.] Bur. Fish., Rep. U.S. Comm. Fish., 1924, append. 8 (Doc. 981):459-486.

LEHMAN, BURTON A.

1953. Fecundity of Hudson River shad. U.S. Fish Wildl. Serv., Res. Rep. 33, 8 p.

LINDSEY, C. C.

1957. The bearing of experimental meristic studies on racial analyses of fish populations. Proc. Ninth Pac. Sci. Congr. 10:54-58.

NICHOLS, PAUL R.

1960. Homing tendency of American shad, Alosa sapidissima, in the York River, Virginia. Chesapeake Sci. 1(3-4):200-201.

RANEY, EDWARD C., and WILLIAM S. WOOLCOTT.

1955. Races of the striped bass, Roccus saxatilis (Walbaum), in southeastern United States. J. Wildl. Manage. 19(4):444-450.

SNEDECOR, GEORGE W.

1956. Statistical methods applied to experiments in agriculture and biology (5th ed.). Iowa State Coll. Press, Ames, Iowa, 534 p.

STEEL, ROBERT G. D., and JAMES H. TORRIE.

1960. Principles and procedures of statistics with special reference to the biological sciences. McGraw-Hill Book Company, Inc., New York, p. 119-123.

TALBOT, GERALD B., and JAMES E. SYKES.

1958. Atlantic coast migration of American shad. U.S. Fish Wildl. Serv., Fish. Bull. 58:473-490.

TÄNING, A. VEDEL.

1952. Experimental study of meristic characters in fishes. Biol. Rev. 27:169-193.

VLADYKOV, V. D., and D. H. WALLACE.

1938. Remarks on populations of the shad (Alosa sapidissima) along the Atlantic coast region. Trans. Amer. Fish. Soc. 67:52-66.

MS #1455

Created in 1849, the Department of the Interior--a department of conservation--is concerned with the management, conservation, and development of the Nation's water, fish, wildlife, mineral, forest, and park and recreational resources. It also has major responsibilities for Indian and Territorial affairs.

As the Nation's principal conservation agency, the Department works to assure that nonrenewable resources are developed and used wisely, that park and recreational resources are conserved for the future, and that renewable resources make their full contribution to the progress, prosperity, and security of the United States--now and in the future.



UNITED STATES
DEPARTMENT OF THE INTERIOR
FISH AND WILDLIFE SERVICE
BUREAU OF COMMERCIAL FISHERIES
WASHINGTON, D.C. 20240

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF THE INTERIOR

OFFICIAL BUSINESS

Return this sheet to above address, if you do
NOT wish to receive this material ☐, or if
change of address is needed ☐ (indicate
change).

~~Dr. I. E. Wallen, Asst. Dir.
for Oceanography
Museum of Natural History
Smithsonian Institute
SSR 7 Washington 25, D. C.~~

